

MARKING SCHEME


SAMPLE PAPER 1


SECTION A



Q.No.	Value Point	Marks
1(i)	D	1
(ii)	B OR A	1
(iii)	B	1
(iv)	C	1
2(i)	B	1
(ii)	A	1
(iii)	A	1
(iv)	A or B	1
3	C	1
4	D or C	1
5	C	1
6	B OR B	1
7	B OR D	1
8	A OR A	1
9	C	1
10	A	1
11	A	1
12	A	1
13	D	1
14	B OR B	1
15	B	1
16	A	1



SECTION B, C, D

Q.No.	VALUE POINTS	MARKS
SECTION B		
17	<p>Nitro group at ortho position withdraws the electron density from the benzene ring and thus facilitates the attack of the nucleophile on haloarene.</p> <p align="center">OR</p>	2

	<p>(i) </p> <p>(ii) $\text{CH}_3\text{CH}(\text{Br})\text{CH}_3 \xrightarrow{\text{alc KOH}} \text{CH}_3\text{CH}=\text{CH}_2 \xrightarrow{\text{HBr, organic peroxide}} \text{CH}_3\text{CH}_2\text{CH}_3\text{Br}$</p>	<p>1</p> <p>1</p>
18	$\Delta T_b = K_f m$ $\Delta T_b = 101.04 - 100 = 1.04^\circ\text{C}$ or $m = 1.04 / 0.52 = 2$ Relative lowering of VP = $x/2$ Relative lowering of VP = $n_2/n_1 + n_2$ $= 2 / 2 + 55.5 = 2 / 57.5 = 0.034 \text{ atm}$	<p>1</p> <p>1/2</p> <p>1/2</p>
19	<p>(i) $t_{2g}^4 e_g^2$ Paramagnetic (ii) Dichloridobis(ethane-1,2-diamine)cobalt(III)nitrate OR (i) Square planar (ii) $\text{Cu}^{2+} = 3d^9$ 1 unpaired electron so $\sqrt{1(3)} = 1.73 \text{ BM}$</p>	<p>1/2, 1/2</p> <p>1</p> <p>1</p> <p>1</p>
20	<p>Reaction is a complex reaction. Order of reaction is 1.5. Molecularity cannot be 1.5, it has no meaning for this reaction. The reaction occurs in steps, so it is a complex reaction. (ii) units of k are $\text{mol}^{-1/2} \text{L}^{1/2} \text{s}^{-1}$ OR Ans : let the rate law expression be $\text{Rate} = k [\text{P}]^x [\text{Q}]^y$ from the table we know that Rate 1 = $3.0 \times 10^{-4} = k (0.10)^x (0.10)^y$ Rate 2 = $9.0 \times 10^{-4} = k (0.30)^x (0.30)^y$ Rate 3 = $3.0 \times 10^{-4} = k (0.10)^x (0.30)^y$ Rate 1 / Rate 3 = $(1/3)^y$ or $1 = (1/3)^y$ So $y = 0$ Rate 2 / Rate 3 = $(3)^x$ or $3 = (3)^x$ So $x = 1$ Rate = $k [\text{P}]$</p>	<p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>1/2</p> <p>1</p>
21	$k = 0.693/t_{1/2}$ $k = 0.693/5730 \text{ years}^{-1}$ $t = \frac{2.303 \log C_0}{k C_t}$ let $C_0 = 1$ $C_t = 3/10$ so $C_0/C_t = 1 / (3/10) = 10/3$ $t = \frac{2.303 \times 5730 \log \frac{10}{3}}{0.693}$ $t = 19042 \times (1 - 0.4771) = 9957 \text{ years}$	<p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1</p>


22	$\begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH} - \text{CH}_3 \\ \quad \\ \text{CH}_3 \quad \text{OH} \end{array} \xrightarrow{\text{H}^+} \begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH} - \text{CH}_3 \\ \quad \\ \text{CH}_3 \quad \text{OH}_2^+ \end{array}$ $\begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH} - \text{CH}_3 \\ \quad \\ \text{CH}_3 \quad \text{OH}_2^+ \end{array} \xrightarrow{-\text{H}_2\text{O}} \begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH}^+ - \text{CH}_3 \\ \\ \text{CH}_3 \end{array}$ $\begin{array}{c} \text{H} \\ \\ \text{CH}_3 - \text{C} - \text{CH} - \text{CH}_3 \\ \\ \text{CH}_3 \end{array} \xrightarrow{1,2\text{-hydride shift}} \begin{array}{c} \text{CH}_3 - \text{C}^+ - \text{CH}_2 - \text{CH}_3 \\ \\ \text{CH}_3 \end{array}$ $\begin{array}{c} \text{CH}_3 - \text{C}^+ - \text{CH}_2 - \text{CH}_3 \\ \\ \text{CH}_3 \end{array} + \text{Br}^- \longrightarrow \begin{array}{c} \text{Br} \\ \\ \text{CH}_3 - \text{C} - \text{CH}_2 - \text{CH}_3 \\ \\ \text{CH}_3 \end{array}$	<p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p>
23	<p>XeF_6</p> <p>. Central atom Xe has 8 valence electrons, it forms 6 bonds with F and has 1 lone pair. According to VSEPR theory, presence of 6 bp and 1 lp results in distorted octahedral geometry</p> 	<p>1</p> <p>1</p>
24.	<p>(a)inverted product will be given by 1 Chlorobutane as it undergoes S_{N}^2 reaction.</p> <p>(b)racemic mixture will be given by 2 chloro-2-methylpropane as it undergoes S_{N}^1 reaction</p>	<p>1/2+1/2</p> <p>1/2+1/2</p>
25	<p>Let no. of Atoms of element P be x</p> <p>No. of tetrahedral voids = $2x$</p> <p>No. Of octahedral voids = x</p> <p>Atoms of Q = $1/3 (2x) + x = 5x/3$</p> <p>$\text{P}_x\text{Q}_{5x/3}$</p> <p>$\text{P}_3\text{Q}_5$</p>	<p>1/2</p> <p>1/2</p> <p>1</p>

SECTION C		
26	<p>(i) Due to large surface area and ability to show variable oxidation states</p> <p>(ii) Due to high value of third ionisation enthalpy</p> <p>(iii) Oxidation state of Cr in Cr₂O₃ is +3 and of CrO is +2. When oxidation number of a metal increases, ionic character decreases so CrO is basic while Cr₂O₃ is amphoteric.</p> <p style="text-align: center;">OR</p> <p>(i) The general trend towards less negative <i>E</i> V values across the series is related to the general increase in the sum of the first and second ionisation enthalpies.</p> <p>(ii) The high energy to transform Cu(s) to Cu²⁺(aq) is not balanced by its hydration enthalpy.</p> <p>(iii) The stability of the half-filled <i>d</i> sub-shell in Mn²⁺ and the completely filled <i>d</i>¹⁰ configuration in Zn²⁺ are related to their more negative <i>E</i>^o V values</p>	1 1 1 1 1 1
27	<p>(i) Aniline, <i>N</i>-ethylethanamine Etanamine</p> <p>(ii) Ethanamine, ethanol, ethanoic acid</p> <p>(iii) <i>N,N</i> dimethylmethanamine, methanamine, <i>N</i>-methylmethanamine</p> <p>OR</p> <p>(i) <i>N</i>-methylethanamine is a secondary amine. When it reacts with benzenesulphonyl chloride, it forms <i>N</i>-Ethyl-<i>N</i> methyl sulphonamide while and <i>N,N</i>-dimethyl ethanamine is a tertiary amine it does not react with benzenesulphonyl chloride.</p> <p>(ii)</p>  <p>(iii) Butan-1-ol</p> <p>Alcohol forms stronger hydrogen bonds with water than formed by amine due to higher electronegativity of O in alcohol than N in amine</p>	1 1 1 1 1 1/2 1/2
28	<p>We know that $d = zM / N_a a^3$</p> <p>For fcc, $z=4$ therefore $d = 4 \times M / N_a (3.5 \times 10^{-8})^3 \text{ g/cm}^3$</p> <p>For bcc, $z=2$ therefore $d' = 2 \times M / N_a (3.0 \times 10^{-8})^3 \text{ g/cm}^3$</p> <p>$d/d' = 4/(3.5 \times 10^{-8})^3 / 2/(3.0 \times 10^{-8})^3 = 3.17:1$</p>	1/2 1 1 1/2
29	<p>(i)</p> 	1


	$\begin{array}{ccccccc} & \text{CH}_2\text{COOH} & & \text{CH}_3 & & (\text{CH}_2)_4\text{-NH}_2 & \\ & & & & & & \\ \text{HOOC} & \text{---CH---} & \text{NHOC---} & \text{CH---} & \text{NHOC---} & \text{CH---} & \text{NH}_2 \end{array}$ <p>(ii)</p> 	1
30	<p>i. Arrange the following in decreasing order of bond dissociation enthalpy $\text{I}_2 < \text{F}_2 < \text{Br}_2 < \text{Cl}_2$,</p> <p>ii. Bi does not form $p\pi$-$p\pi$ bonds as its atomic orbitals are large and diffuse so effective overlapping is not possible</p> <p>iii. Due to small size of oxygen, it has greater electron electron repulsions</p>	1 1 1
SECTION D		
31.	<p>(i)</p> <p>(a) $3\text{Cu} + 8\text{HNO}_3(\text{dilute}) \rightarrow 3\text{Cu}(\text{NO}_3)_2 + 2\text{NO} + 4\text{H}_2\text{O}$</p> <p>(b)</p>  <p>(ii) 'X' is Helium It is used as a diluent for oxygen in modern diving apparatus because of its very low solubility in blood. It monoatomic having no interatomic forces except weak dispersion forces and has second lowest mass therefore bp is lowest.</p> <p style="text-align: center;">OR</p> <p>(a) $\text{H}_2\text{Te}, \text{H}_2\text{Se}, \text{H}_2\text{S}, \text{H}_2\text{O}$</p> <p>(b) $[\text{Fe}(\text{H}_2\text{O})_5(\text{NO})]^{2+}$</p> <p>(ii) A is chlorine gas Its bleaching action is due to oxidation. $\text{Cl}_2 + \text{H}_2\text{O} \rightarrow 2\text{HCl} + \text{O}$, Coloured substance + O \rightarrow Colourless substance $6\text{NaOH} + 3\text{Cl}_2 \rightarrow 5\text{NaCl} + \text{NaClO}_3 + 3\text{H}_2\text{O}$</p>	1 1 1 1 1 1 1 1 1
36		

½ each


A: $\text{C} \equiv \text{CH}$




B: COCH_3



C: COOK D: CHI_3




$\text{C} \equiv \text{CH}$




$\xrightarrow{\text{HgSO}_4, \text{H}_2\text{SO}_4}$

COCH_3




1



$\xrightarrow{\text{CH}_3\text{COCl, anhy AlCl}_3}$


COCH_3



1


1

COCH_3



$\xrightarrow{\text{KOH, I}_2}$

COOK



+ CHI_3

OR

(i) $\text{CH}_3\text{COCH}_3 + \text{CH}_3\text{CHO}$

\downarrow dil NaOH


$(\text{CH}_3)_2\text{C}(\text{OH})\text{CH}_2\text{CHO} + \text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{COCH}_3$

\downarrow Heat

$(\text{CH}_3)_2\text{C}=\text{CHCHO} + \text{CH}_3\text{CH}=\text{CHCOCH}_3$

1

1



$$A = -\text{slope} = - (149 - 147.8 / 0.010 - 0.022) = 100 \text{ S cm}^2 \text{ mol}^{-1} / (\text{mol/L}^{-1})^{1/2}.$$